

Results of the Profile Gravity Measurements in the Pieniny Klippen Belt (Middle Váh Valley, Western Slovakia)

Jana Bučová¹, Miroslav Bielik², Roman Pašteka², Dušan Plašienka³, Roman Aubrecht^{1,3}, Juraj Papčo⁴

¹ Geophysical Institute, Slovak Academy of Sciences, Dúbravská cesta 9, 845 04 Bratislava, Slovakia;
geofjabu@savba.sk

² Department of Applied and Environmental Geophysics, Faculty of Natural Sciences, Comenius University, Mlynská dolina, 842 15 Bratislava 4, Slovakia; *bielik@fns.uniba.sk, pasteka@fns.uniba.sk*

³ Department of Geology and Paleontology, Faculty of Natural Sciences, Comenius University, Mlynská dolina, 842 15 Bratislava 4, Slovakia; *plasienska@fns.uniba.sk, aubrecht@fns.uniba.sk*

⁴ Department of Theoretical Geodesy, Faculty of Civil Engineering, Slovak University of Technology, Radlinského 11, 813 68 Bratislava, Slovakia; *juraj.papco@stuba.sk*

The examined area is situated in the Púchov section of the Pieniny Klippen Belt (PKB) in western Slovakia, on the territory of municipalities Tuchyňa, Mikušovce, Červený Kameň and Kvašov. The main aim of our research was to broaden the present knowledge on geological structure of the PKB and on its tectonic connection to the surrounding regional units using the geophysical methods.

The profile gravity measurements consist of two parallel profiles named *prof_1* and *prof_2* of roughly NNW-SSE direction. The eastern section *prof_1* has a length of 6.36 km; the western profile *prof_2* is about 3.84 km long. Measurements were carried out at intervals of 20 meters (overall 506 points were gauged) by gravimeters Scintrex CG3 and CG5. The acquired data were processed into the Bouguer gravity anomalies. The software package GM-SYS (Geosoft) was applied for the gravity modelling and the final geological-geophysical sections represent distribution of the lithological members with their specific density parameters (density values were determined by the own laboratory measurements of rock samples collected along the profiles). The density modelling was also based on the results obtained by the geoelectrical exploration (vertical electric sounding and resistivity profiling), new surface geological mapping and borehole interpretation (continuously cored bores MIK-1 and MIK-2).

Presented detailed gravity measurements reflect density differences between klippen (rigid limestone blocks) and their mantle (predominant marlstone) and they demonstrate that the primary fold-nappe system of the PKB was destructed by the subsequent deformation, resulting in the formation of the positive flower structure and backthrusting. Summing up, the new density modelling confirmed complex geological structure of the studied area and helped us to better interpret the tectonic evolution of the PKB.

Acknowledgements: The authors are thankful to the Slovak Research and Development Agency (grants APVV-0465-06, APVV-0212-12 and LPP-0225-06) and the Slovak Grant Agency VEGA (grant No. 2/0195/12) for the financial support.

* * *